

Nutritionally complete, low fat, powdered formula, low in long-chain triglycerides (LCT) and high in medium-chain triglycerides (MCT). Lipano contains a dosed amount of vitamins, minerals, trace elements, docosahexaenoic acid (DHA) and arachidonic acid (AA).



Product USPs

Contains low LCT content and high MCT content

Matched to requirements with carbohydrates, vitamins, minerals, trace elements, and omega-3 and omega-6 essential fatty acids

Definition and Indication:

FOOD FOR SPECIAL MEDICAL PURPOSES

For the dietary management of fat maldigestion and fat malabsorption, oxidation disorders of long chain fatty acids and in all other cases that require an additional source of MCT fats.

Ingredients:

maltodextrin, vegetable oils 16,6% (medium chain triglyceride from palm and coconut oil in varying proportions, **soybean** oil, antioxidants: tocopherol rich natural extract, ascorbil palmitate), calcium caseinate (**milk**), lactose (**milk**), fructo-oligosaccharides (FOS) 3,4%, di-Calcium phosphate, tri-Potassium citrate, emulsifiers: **soya** lecithin, citric acid esters of mono- and diglycerides of fatty acids; galactooligosaccharides - GOS (**milk**) 0,5%, sodium hydrogen phosphate, potassium chloride, choline bitartrate, magnesium carbonate, vitamin (A, D, E, K, C, B1, B2, niacin, B6, folic acid, pantothenic acid, B12, biotine), sodium chloride, oil from *Mortierella alpina*, DHA- and EPA-rich oil derived from the micro alga *Schizochytrium sp.*, L-cystine, inositol, iron sulphate, zinc sulphate, L-carnitine L-tartrate, copper sulphate, manganese sulphate, potassium iodide, sodium selenite, sodium molybdate.

Important notices

Use under medical supervision.

Not for parenteral use.

Not to be used by individuals who do not have the specific disease.

Suitable for the use as a sole source of nutrition in infants or as a supplement for children over the age of one and adults.

Dosage

to be determined by the clinician or dietitian and is dependent on the age, body weight and medical condition of the patient.

Contraindications

Lipano should not be used in cases of suspected ketoacidosis/acidosis, decompensated liver cirrhosis, chronic renal insufficiency or in patients with medium chain acyl-CoA dehydrogenase deficiency (MCADD). patients with medium chain acyl-CoA dehydrogenase deficiency (MCAD)

Preparation guidelines

Lipano can be administered orally or via a feeding tube, according to the instructions of the doctor or nutritionist. Standard dilution (15%): the recommended concentration is 90 ml of water plus 15 g of Lipano.

Storage

Closed: Store at room temperature in a dry place. Opened: Once opened, remember to put the lid on and consume within 4 weeks.

Nutritional Declaration

Average values for:	100 g	per Portion*
Energy kJ	1.929 kJ	1.929 kJ
Energy kcal	460 kcal	460 kcal
Fat	17,5 g	17,5 g
of which saturates	14,3 g	14,3 g
of which mono-unsaturates	0,7 g	0,7 g
of which poly-unsaturates	2,4 g	2,4 g
of which MCT	13,4 g	13,4 g
of which LCT	4,5 g	4,5 g
of which linoleic acid	1,7 g	1,7 g
of which α-linolenic acid (Omega 3)	0,3 g	0,326 g
of which arachidonic acid (ARA)	70,0 mg	70,0 mg
of which docosahexaenoic acid (DHA)	56,0 mg	56,0 mg
Carbohydrates	63 g	63 g
of which sugar	8,1 g	8,1 g
Fibre	3,4 g	3,4 g
Protein	10,9 g	10,9 g
Salt	0,32 g	0,32 g
of which sodium	129 mg	129 mg
Vitamins		
Vitamin A (RE)	429 µg	429 µg
Vitamin D	12 µg	12 µg
Vitamin E (α-tocopherol)	4,29 mg	4,29 mg
Vitamin K	18 µg	18 µg

Vitamin C	69 mg	69 mg
Thiamin B1	343 µg	343 µg
Riboflavin B2	686 µg	686 µg
Niacin B3	3,50 mg	3,50 mg
Vitamin B6	429 µg	429 µg
Folate B9 (DFE)	129 µg	129 µg
Folic acid B9	77 µg	77 µg
Vitamin B12	0,86 µg	0,86 µg
Biotin B8	8,58 µg	8,58 µg
Pantothenic acid B5	3,00 mg	3,00 mg
Minerals		
Potassium	386 mg	386 mg
Chloride	283 mg	283 mg
Calcium	429 mg	429 mg
Phosphorus	344 mg	344 mg
Magnesium	64 mg	64 mg
Trace Elements		
Iron	4,88 mg	4,88 mg
Zinc	4,29 mg	4,29 mg
Copper	300 µg	300 µg
Manganese	214 µg	214 µg
Selenium	15 µg	15 µg
Molybdenum	11 µg	11 µg
Iodine	73 µg	73 µg
Others		
L-Carnitine	6,43 mg	6,43 mg
Choline	132 mg	132 mg
Inositol	53 mg	53 mg

Osmolality 130 mOsm/kg

*One portion is equivalent to 15g. Standard dilution: 15g Lipano (3 spoons) + 90 ml drinking water = 100 ml ready-to-use feed.

Order information

Send an e-mail to info@kanso.com to learn if this product is available in your country.

OUTER CARTON

Units per carton	6
Net Gross weight	3.000 g 3.889 g
Dimensions (H x W x D)	17 x 33 x 22.6 cm

PALLET

Units per pallet	432
Cartons per pallet	72
Cartons per layer	12
Layer per Pallet	6
Pallet dimensions in cm (height x width x depth)	116.5 x 80 x 120 cm
Gross weight	301 kg

SINGLE PACK

Item Number	6069080001
EAN Unit	8008698032022
Shelf life	540 days
Net Gross weight	500,0 g 620,0 g
Dimensions (H x W x D)	14,9 x 10,3 x 10,3 cm



This datasheet is subject to change. Please consult the product packaging for legally binding information about individual products.